0 - 31865 - 30924 - 3

Visit our website: www.AndersonsPlantNutrient.com

©2019 The Andersons, Inc. The Andersons logo is a registered trademark of The Andersons, Inc.; Professional Turf is a trademark of The Andersons, Inc.; SURGE is a registered trademark of PBI/Gordon Corporation 082619

> **APT164SR4.6 APT164SR4.6B**

## The Professional Andersons Turf Products

## Fertilizer with SURGE®

Net Wt. 40 Lbs. (18.14 kg)

Broadleaf Herbicide Not for sale, distribution or use in Nassau or Suffolk Counties in New York State.

## 16-0-9

This product is a herbicide & plant nutrient combination which controls a wide spectrum of broadleaf weeds while providing both fast & long-lasting feeding action.

Combination of 4 active ingredients enhances speed of herbicide Superior cool season performance.

· Controls tough broadleaf weeds. Mecoprop-p acid . .

OTHER INGREDIENTS: THIS PRODUCT CONTAINS: 15.30 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.765% (CAS 1928-43-4) 3.48 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.174% 1.46 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.073% (CAS 1918-00-9) 0.68 lb N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]

EPA Reg. No. 2217-882-9198 EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A Underlined letter is first letter used in run code on bag **GUARANTEED ANALYSIS** 

phenyl]methanesulfonamide per ton or 0.034% (CAS 122836-35-5)

Soluble Potash (K2O) . . . Derived from urea, potassium chloride and polymer coated sulfur coated urea. \*4.00% Slowly available nitrogen from polymer coated sulfur coated urea. Information regarding the contents and levels of metals in this product is available on the

> KEEP OUT OF REACH OF CHILDREN CAUTION STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals CAUTION: Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes or

Personal Protective Equipment (PPE) All loaders, applicators, and other handlers must wear:

internet at http://www.regulatory-info-ap.com.

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

**User Safety Recommendations** 

• Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the • Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean

If in eyes: • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue Call a poison control center or doctor for treatment advice.

 Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or Do not give anything by mouth to an unconscious person. Have the product container or label with you when calling a poison control center or doctor or

going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment

swallowed: • Call a poison control center or doctor immediately for treatment advice.

**ENVIRONMENTAL HAZARDS** This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of

equipment wash waters or rinsate. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

• Do not enter or allow people (or pets) to enter the treated area until dusts have settled. Do not apply this product in a way that will contact workers or other persons, either directly or • Not for use on turf being grown for sale or other commercial use as sod, or for commercial

 Do not use this product on dichondra, carpetgrass, creeping bentgrass, or turf containing desirable clovers. • Do not apply this product to newly seeded grasses until well-established or after the third mowing. Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area. Do not use this product on 'Floratam' St. Augustinegrass.

• Do not use this product on or near desirable plants, including within the drip line of the roots of

desirable trees and shrubs, since injury may result. This product is a Surge® Broadleaf Herbicide, containing fertilizer, 2,4-D, mecoprop-p, dicamba and sulfentrazone. This granular weed and feed provides plant nutrients and controls a broad spectrum of listed broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorus, and potash

Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the should be applied according to soil tests or State Extension Service recommendations. This product is intended to be applied by noncommercial users, lawn care operators, and

landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated. Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations; for example: hospitals, nursing homes, schools, museums, libraries, sport facilities (including golf courses), and office buildings may be treated. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings. Finally, residential sites are defined as areas associated with the ousehold or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes. his product may be applied to newly seeded cool season turfgrass. Apply only after the third mowing or after the grass seedlings have tillered and have developed an adequate secondary

This product may be applied to newly established warm season turfgrass. Treatments should be providing 0.32 - 0.64 pounds of nitrogen per 1,000 square feet and herbicide to control listed delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations, and low application rates (1/2 of the listed rate for established turfgrass) are suggested.

A uniform broadcast distribution of product on the treated area is essential. Do not exceed the specified application rate for any area. Do not apply product to bare ground. Do not use this product on dichondra, carpetgrass, creeping bentgrass, or turf containing desirable clovers. California: Make broadcast applications only between March 1 and September 1. While irrigation is necessary and important for plant growth, apply irrigation water efficiently so that no more than 125% of the net irrigation requirement is applied for any irrigation event. Apply efficient irrigations for six months following application of sulfentrazone containing products. **New York**: Only one application per year of this product is allowed. This product is not allowed to

HOW MUCH TO APPLY:
For Kentucky bluegrass, fescue sp., pe en gill rye r.s. om non bermudagrass, and zoysiagrass: Apply 130-174 pounds of this product per acre of established turfgrass. The equivalent application rates are 3.0 - 4.0 pounds of product per 1,000 square feet of turfgrass. For New York State only: Apply 130-166 pounds of this product per acre of established turfgrass. The equivalent application rates are 3.0 - 3.8 pounds of product per 1,000 square feet

For St. Augustinegrass, centipedegrass, hybrid bermudagrass, bentgrass, and colonial **bentgrass:** Apply 87-109 pounds of this product per acre of established turfgrass. The

equivalent application rates are 2.0 - 2.5 pounds of product per 1,000 square feet of turfgrass. imitations on broadcast applications to ornamental turfgrass. Maximum Rate | Maximum Number | Minimum Inter- | Maximum Seasonal Use Site per Application of Applications val Between

Turfgrass 174 pounds/A 30 Days Do not exceed the maximum seasonal rate of 348 pounds, per acre per season, excluding spot Sequential or follow-up applications may be required for dense infestations of perennial and

biennial weeds. Also, adverse environmental conditions prior or after application may reduce the effectiveness. If required, the treatment interval between the first and second applications should be approximately 30 days. Drop spreaders, rotary spreaders, and other granular equipment should be calibrated for accurate and uniform applications of this product. Calibrate the equipment by applying the product over a measured or known area and adjust the equipment according to the recommendations of the equipment manufacturer. Refer to the rate table for the appropriate bag sizes and

APPLICATION DIRECTIONS:

. Spring and fall applications are suitable for the best control. However, this product can be applied any time broadleaf weeds are present and actively growing. 2. Mow the turfgrass 1 to 2 days before application. 3. Water the turfgrass thoroughly (>1/2 inch) at least 1 to 2 days before application. I. Schedule applications in the morning when dew is present. Or, sprinkle or irrigate lightly (<1/4 inch) before application to provide maximum control. 5. Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.

> SPREADER SETTINGS These suggested spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product. 40 lbs. treats 16,000 sq. ft. at the 109 lbs/acre - LOW RATE (2.5 lbs/1,000 sq. ft.) 40 lbs. treats 10,500 sq. ft. at the 166 lbs/acre - MEDIUM RATE (3.8 lbs/1,000 sq. ft.) 40 lbs. treats 10,000 sq. ft. at the 174 lbs/acre - HIGH RATE (4.0 lbs/1,000 sq. ft.) GROUND | WIDTH OF | SPREADER SETTINGS

	SPEED	COVERAGE			
SPREADER			LOW RATE	MEDIUM RATE	HIGH RATE
Andersons Model 2000/AccuPro 2000 SR/SR-2000 (cone 6)	3 mph	8 feet	K	L 3/4	М
Scotts R-8A (cone 6)	3 mph	8 feet	J	K 3/4	K 1/2
LESCO 020093 (.25)	3 mph	6 feet	D	Е	E 1/4
LESCO 705698 (.25)	3 mph	6 feet	10	12	12 1/2
Lely Models WTR, WFR, HR, 1250 (PTO at 450 rpm)	4.5 mph	26 feet	4 II-C	5 II-C	5 1/4 II-C
Vicon (03 series)	4.5 mph	20 feet	19	24	25

To provide proper distribution, calibrate spreader before application. For 2.5 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 400 square feet and adjust to apply at the rate of 1 pound of product per 400 square feet. For 3.8 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 260 square feet and adjust to apply at the rate of 1 pound of product per 260 square feet. For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

granules evenly. Never apply by hand. Coverage - To determine how much you need to apply, multiply the length of your lawn by its

to change directions or when stopping.

width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One 40 pound bag covers up to 20,000 square feet. SINGLE PASS DISTRIBUTION: For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps

1. For rotary spreaders, operate at intervals corresponding to the swath width. For drop spreaders, overlap the wheel tracks to ensure complete coverage. 2. First, spread the borders of the lawn. Then apply this product to the center with straight line 3. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. weeds. Please consult your local extension office for questions regarding a regular fertilizer

Dandelion, chickweed, plantain, and clover are controlled effectively with this product. Mouseear hawkweed Strawberry, India mod Parslev-piert Tansy ragwort Tansy mustard Trailing crownvetch Venice mallow Virginia buttonweed Wavvleaf bullthistle Western clematis Prostrate knotweed Wild four-o'-clock Prostrate pigweed Wild geranium Wild lettuce Purslane, common Redroot pigweed Rough cinquefoil Russian pigweed Sheep sorrel Woolly plantain Shepherdspurse Slender plantain Smallflower galinsoga Yellow rocket Smooth dock Yellowflower pepperwee

## STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage and disposal **PESTICIDE STORAGE:** Store in original container and keep any bags containing unused uct in a locked storage area inaccessible to children and domestic animals. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law, If these wastes cannot be disposed of b according to label instructions, contact your State Pesticide or Environmental Control ency, or the Hazardous Waste representative at the nearest EPA Regional Office for

**CONTAINER HANDLING:** Nonrefillable container: Do not reuse or refill this container. ompletely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER
FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW,
THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED

WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTUR-ER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULT-ING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

> Maumee, Ohio 43537 **SGN 150**

The Care Professional Andersons Turf™ Products

Fertilizer with SURGE® **Broadleaf Herbicide** 

NO PRINT NO COPY

Wit

NO PRINT NO COPY