

Net Wt. 50 Lbs. (22.68 kg)
This product contains 0.075 lbs. of Pendimethalin per lb. of product.

22-0-6 Fertilizer with ProPendi™ Herbicide

 Preemergent control of crabge goosegrass, foxtail, chickweed oxalis, spurge, knotweed, and annua. 	l, É
 Contains NS-54™ for controlle nitrogen feeding. CTIVE INGREDIENT: 	d release
endimethalin†	0.75%
VERT INGREDIENTS:	
	al 100.00%
CAS #40487-42-1	
PA Reg. No. 9198-177	
PA Est. No. 538-0H-1S, 9198-0H-1M, 9198-0H-2B, 9198-AL-0	01A
Inderlined letter is first letter used in run code on bag.	_
juaranteed Analysis	
otal Nitrogen (N)	22.00%
22.00% Urea Nitrogen*	
oluble Potash (K,0)	6.00%

*8.80% Slowly available urea nitrogen from polymer coated sulfur coated

Information regarding the contents and levels of metals in this product is available on the internet at http://www.regulatory-info-ap.com We recommend that you follow the Green Industries BMP's at: http://fyn.if-

as.ufl.edu/pdf/GIBMP_Manual_Web_English_2015.pdf We recommend that you follow the Golf Course BMP's at: http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/glfbmp07.pdf **Nitrogen Notice:** For those states with nitrogen application restrictions, including but not limited to, MD, NH, NJ and VA, do not apply more than 3.83 lbs of this product per 1,000 sq.ft. U.S. Patent No's 3,705,794; 3,989,470

*Licensed under one or more of the following U.S. Patents: 3,920,742;

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals: CAUTION, causes eye irritation. Avoid contact with eyes, skin, or clothing. In case of contact, immediately flush eyes or skin

KEEP OUT OF REACH OF CHILDREN

with plenty of water. Get medical attention if irritation persists. May be harmful if treated areas. Do not feed clippings to livestock. **If swallowed:** Call a poison control center immediately for treatment advice. Have

person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an **If on skin:** Take off contaminated clothing. Rinse skin immediately with plenty of

water for 15-20 minutes. Call a poison control center or doctor for treatment advice. If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling poison control center or doctor, or going for treatment. You may contact 1-800-757-8951 for emergency medical treatment information. Personal Protective Equipment

Applicators and other handlers must wear: Long-sleeved shirt and long pants, chemical resistant gloves, made of waterproof materials, such as butyl rubber >14 mils, nitrile rubber > 14 mils or neoprene rubber > 14 mils, shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining personal protection equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash personal protection equipment separately from other laundry.

 Wash hands before eating, drinking, chewing gum, using tobacco or using the put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Engineering Controls:

the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS. This pesticide is toxic to fish. Do not apply directly to water, to areas where surface

either collect for use or dispose of properly. Non-target organism advisory statement: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including

pollinators, in areas adiacent to the treated site. Protect the forage and habitat of

non-target organisms by following label directions intended to minimize spray drift.

sistant biotypes may dominate the weed population if these herbicides are used sprout.

(1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of proportional department of the proportional non-controlled plants of a particular weed species; (3) surviving plants mixed with

F002091 with a different mode of action, if available. recommendations for specific types of plants and weed biotypes.

AGRICULTURAL USE REQUIREMENTS

gricultural workers on farms, forests, nurseries, and greenhouses and handlers of

Andersons Model 2000/

AccuPro 2000; (cone

4) Andersons 2000 SR/

SR-2000 (cone 4)

Lely Models WTR, WFR,

• Rotate the use of this product or other Group KI herbicides within a growing season

Where lower infestations are expected, the lighter rates may be used. sequence or among growing seasons with different herbicide groups that control the same

If a weed pest population continues to progress after treatment with this product, turfgrass is under severe stress conditions (disease, drought, low temperature - below 40°F, scalping, Comments discontinue use of this product and switch to another management strategy or herbicide

and excessive soil moisture).

Apply a repeat application of 167-200 lbs./acre after 6-8 weeks for extended control or where heavy week infestations are expected. Contact your sales representative, pest control advisors, or local extension specialist for additional pesticide resistance management and/or integrated weed management

andard 40 CFR part 170. This standard contains requirements for the protection of ricultural pesticides. It contains requirements for training, decontamination, notificatio d emergency assistance. It also contains specific instructions and exceptions pertaining

tandard and that involves contact with anything that has been treated, such as plants, soil or water is: Long-sleeved shirt and long pants, chemical resistant gloves, made of waterproof

Seeding or Sprigging Intervals after Treatment materials, such as butyl rubber > 14 mils, nitrile rubber > 14 mils, or neoprene rubber > 14 Delay seeding by 3 months and sprigging by 5 months after application. Prior to planting, for best

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter Do not apply more than a cumulative total of 400 lbs. of product/acre (3.0 lbs. active ingredient/acre) treatment areas until spray has dried or dusts have settled.

per year on bentgrass and *Poa annua*; 800 lbs. of product/acre (6.0 lbs. active ingredient/acre) per year For use on any turfgrass site (lawns, sod, turf areas) including, but not limited to: grounds or lawns on cool season turfgrass; and 1,200 lbs. of product/acre (9.0 lbs. active ingredient/acre) per year on around residential and commercial establishments, golf courses, parks, athletic fields, schools, warm season turfgrasses.

For use on established turfgrass including bentgrass, bermudagrass, and *Poa annua* (except putting n of Federal law to use this product in a manner inconsistent with its labeling. greens and tees), Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, St. Augustinegrass, bahiagrass, zoysiagrass, centipedegrass and blends of the grasses. See "Bermudagrass, Bentgrass and this product in a way that will contact workers or other persons, either directly Poa annua Precautions" and "Limitation of Liability" statements. ements specific to your State or Tribe, consult the agency responsible for pesticide Not for use on bentgrass, *Poa annua* and bermudagrass putting greens and tees, or on dichondra. Where bentgrass, bentgrass-containing turf and *Poa annua* are the desired species, a light rate

develop plants naturally resistant to this product and other Group K1 herbicides. fall panicum, Poa annua, oxalis, knotweed, purslane, and prostrate spurge, apply in spring before they

and uses historical information related to herbicide use and that considers mechanical product to the soil surface for an effective chemical barrier. Do not destroy the chemical barrier creations are supported by the chemical barrier creations and uses historical information related to herbicide use and that considers mechanical control methods, cultural (e.g. timing to favor the desirable plants and not the weed), at the soil surface by raking or aerifying as a segment to application. Allow at least a four month sp biological (weed-competitive varieties) and other management practices.

• Scout after herbicide application to monitor weed populations for early signs of resistance

Do not mow treated area until product has been watered in and turf foliage has dried.

Do not mow treated area until product has been watered in and turf foliage has dried.

Social formula (1/2 inch poa Annua height or taller.) Not Barnyardgrass.

Where bentgrass and bentgrass-containing turf and *Poa annua* are the desired species, a light rate controlled individuals of the same species. If resistance is suspected, prevent weed seed application only is recommended. production in the affected area by an alternative herbicide from a different group or a

On cool season turfgrasses avoid applications when temperatures are regularly in the 80's.

with cool season turfgrass species may injure and thin the overseeding.

Bahiagrass

Barnyardgrass

staining, If necessary, rinse with water. or less) Purslane If metallic surfaces (golf course equipment, automobiles, etc.) are present at the application site, do Tall Fescue Knotweed not use this product under windy conditions (5-10 MPH) due to its staining potential. (Not for use on greens Chickweed May cause temporary slowing of turf fill-in and stolen-rhizome rooting where turf has been thinned.

Suggested Spreader Settings

To provide proper distribution calibrate spreader before application.

Do not adjust rates in order to meet fertilizer requirements of your soil.

50 lbs. treats 13,000 sq. ft. (0.30 acre) at the 167 lbs. product/acre - LIGHT RATE (3.83 lbs./1,000 sq. ft.)

50 lbs. treats 10,800 sq. ft. (0.25 acre) at the 200 lbs. product/acre – LOW RATE (4.59 lbs./1,000 sq. ft.)

50 lbs. treats 9,300 sq. ft. (0.21 acre) at the 233 lbs. product/acre - MEDIUM RATE (5.35 lbs./1,000 sq. ft.)

50 lbs. treats 8,100 sq. ft. (0.19 acre) at the 266 lbs. product/acre - HIGH RATE (6.10 lbs./1,000 sq. ft.)

Andersons LCO-1000 3 mph 10 feet L 3/4 M 1/2 N 3/4 0 1/2

Scotts R-8, R-8A (cone 4) 3 mph 10 feet K 3/4 L 1/2 M 3/4 N 1/2

HR, 1250 (PTO at 450 4.5 mph 34 feet 6 6 1/2

ESCO 020093 (0.25) 3 mph 7 feet E E 1/2 F F1/2

Ground Width of LIGHT RATE LOW RATE MEDIUM RATE HIGH RATE

Speed Coverage 1.25 lbs. ai/acre 1.50 lbs. ai/acre 1.75 lbs. ai/acre 2.00 lbs. ai/acre

TURFGRASS SPECIES	WEEDS	RATES
Cool Season Grasses	Crabgrass	200-266 lbs./acre as the
Kentucky bluegrass	Foxtail	initial application prior
Perennial ryegrass	Poa Annua	to weed germination.
Fine fescue	Barnyardgrass	
Tall fescue	Fall Panicum	
	Oxalis	
	Prostrate Spurge	
	Purslane	
	Knotweed	
	Chickweed	

The first of the second for the	ition of 167-200 lbs./acre estations are expected.	after 6-8 weeks for extended control or
	Goosegrass	200-266 lbs./acre in the spring prior t weed germination. Apply a repeat application after 6-8 weeks at 200-233 lbs./acre
Bentgrass and	Crabgrass	200 lbs./acre in the spring prior

heavy weed infestations are expected. Goosegrass 200 lbs./acre in the spring prior to weed

sandy soils.

Wafin season grasses Craugiass 200 200 into george
Bermudagrass Foxtail germination.

Goosegrass 200-266 lbs./acre in the spring prior to Cudweed weed germination with a repeat application of 200-266 lbs./acre

STORAGE AND DISPOSAL Do not contaminate water, food or feed by storage or disposal. **PESTICIDE STORAGE:** Store in a clean, di place. Reseal opened bag by folding top down and securing. PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If not available, then dispose of empty bag in a sanitary landfill or by

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using the product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability: WARRANTY DISCLAIMER and LIMITATION of LIABILITY: Manufacturer warrants that this product conforms to the chemical description on the label and is

reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted. THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY

 Vicon (03 Series)
 4.5 mph
 28 feet
 29
 33
 36
 38
 P.O. Box 119, Maumee, OH 43537

FROM STATE/JURISDICTION TO STATE/JURISDICTION.





0 31865 32687 5

Visit our website at www.andersonsplantnutrient.com

©2021 The Andersons, Inc. The Andersons logo is a registered trademark of The Andersons, Inc. Professional Turf is a trademark of The Andersons, Inc. ProPendi is a trademark of The Scotts Company

APT228PD50.5

The ***
Andersons

22-0-6 Fertilizer with ProPendi™ Herbicide