


Andersons Golf Products

DG Lime™



Dispersing Granule Technology

- **Dispersing granule formula**
- **Provides calcium & magnesium, essential for plant growth**
- **Pelletized for easy application**
- **Neutralizes acidic soils**

Guaranteed Analysis	
Calcium (Ca)	30.00%
Magnesium (Mg)	4.00%
Calcium Carbonate (CaCO ₃).....	75.00%
Calcium Oxide (CaO).....	42.00%
Magnesium Carbonate (MgCO ₃)	13.00%
Magnesium Oxide (MgO)	6.00%
Calcium Carbonate Equivalent (CCE)	90.00%

Derived from dolomite

ALSO CONTAINS NON-PLANT FOOD INGREDIENT:
2% lignosulfonates as Water Soluble Binder

FO02091

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>.

Pennsylvania, Ohio: ENP = 1,800 lbs.
Missouri: ENM = 630 lbs./ton
Iowa: Iowa Secretary of Agriculture Certified 1,500 lbs. ECCE per ton
Florida: This material required 1,858 lbs. to be equal to one ton of standard agricultural liming material.
Kansas, Nebraska, Oklahoma: ECCE = 75%
Minnesota: ENP = 82%
North Carolina: 1,886 lbs. Of this material equals one ton of standard agricultural liming materials ENV = 90%
Tennessee: RNV = 80%
New York: Effective Neutralizing Value = 79.2%; 1.1 tons of material needs to equal one ton of material having 100% Effective Neutralizing Value
Wisconsin: Neutralizing Index Zone = 80-89
Oregon: Lime Score = 95
Virginia: Effective Neutralizing Value = 90%

FLORIDA APPLICATIONS:
We recommend that you follow the Green Industries BMP's at:
http://fyn.ifas.ufl.edu/pdf/GIBMP_Manual_Web_English_2015.pdf
We recommend that you follow the Golf Course BMP's at:
<http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/gfbbmp07.pdf>



WARNING
May be harmful if swallowed
May cause mild skin irritation
May cause eye irritation
May cause respiratory irritation

- Precautionary Statements:**
- Call a poison center/physician if unwell
 - If skin irritation occurs, get medical advice
 - If in eyes rinse cautiously with water for several minutes - remove contact lenses
 - If eye irritation persists, get medical advice
 - Wash hands after handling
 - Use only outdoors or in well ventilated areas
 - Avoid breathing dust
 - If inhaled remove to fresh air and keep at rest in a position comfortable for breathing
 - Store locked up
 - Dispose of contents/container in accordance with national/regional/local regulations

First Aid:
If inhaled: Move victim to fresh air. Seek medical attention if irritation persists.
If on skin: Wash affected areas with soap and water. Seek medical attention if irritation persists. Wash contaminated clothing before re-use.
If in the eyes: Immediately flush with water for at least 20 minutes. Seek medical attention if irritation persists.
If swallowed: If victim is alert and not convulsing, give one glass of water to dilute material. Seek immediate medical attention.

Directions for Use: Apply only as directed. For use on turf and in gardens. The pellets in this product disperse immediately into the soil to reduce soil acidity. When used according to label directions, this product is non-toxic to humans, pets and plant life. For best results, determine the pH levels of the area to be treated by conducting a pH soil test. If that information is not available, refer to the recommendations below on rates and timing.

When establishing a new turf area, apply 40 lbs. per 1,000 sq. ft. prior to seeding. Rake lightly into the soil. Water well after application.

For established turf areas, apply 25-40 lbs. per 1,000 sq. ft. For clay soils, apply 18 lbs. of product per 1,000 sq. ft.

To maintain proper pH soil levels, repeat application 1 to 2 times per year, or when the soil pH level drops below 6.3, at rates of 4-6 lbs. per 1,000 sq. ft.

To correct pH levels, apply 40 lbs. per 1,000 sq. ft. or 2 lbs. per 1,000 sq. ft. for every 1/10 pH correction needed.
Gardens have a deeper root zone and require a higher application rate than turf. For best results, double the turf recommended rates and mix into the soil at a depth of 6 inches.
For all applications: Immediately after application, apply a minimum of 0.5 inches of water to disperse particles into the root zone. Due to variations in temperature and water pH, best results are achieved by splitting the irrigation into 2 applications at 10-15 minutes apart. This will allow for more thorough particle dispersion into the root zone.

For incorporation, after application, wet thoroughly 30 minutes before tilling.

Fineness of Grind
Prior to pelletizing or testing by wet sieve method:

Minimum Passing	U.S. Standard Sieves
100%	8 & 20 MESH
98%	40 MESH
94%	50 MESH
85%	60 MESH
70%	100 MESH

IMPORTANT: READ BEFORE USE:
Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:
Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS.

LEGAL RIGHTS:
THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Product of USA

Distributed by:
The Andersons, Inc.
P. O. Box 119
Maumee, Ohio 43537
www.andersonsplantnutrient.com

DG Lime is a trademark of The Andersons, Inc.

Suggested Spreader Settings						
These suggested spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product						
SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS			
			LIGHT RATE 2 lbs/1,000 sq. ft.	LOW RATE 4 lbs/1,000 sq. ft.	MEDIUM RATE 12 lbs/1,000 sq. ft.	HIGH RATE 20 lbs/1,000 sq. ft.
Andersons Model 2000 / AccuPro 2000 (cone 9); Andersons 2000 SR / SR-2000 (cone 9)	3 mph	10 feet	I	J 1/2	Q	X
Scotts R-8, R-8A (cone 9)	3 mph	10 feet	G 1/2	I	O 1/2	V 1/2



SGN 100
Net Wt. 50 Lbs (22.68 kg)

10004946 (AGCDGLIME5.40)