IMENSION®



Fertilizer with 0.164% DIMENSION® Turf Herbicide

Net Wt. 50 Lbs. (22.68 kg)

in established lawns and ornamental turfs.

18.00% Urea Nitrogei Soluble Potash (K,O). Derived from urea, polymer coated sulfur coated urea and potassium chloride. *3.60% Slowly available urea nitrogen from polymer coated sulfur coated urea.

Nitrogen Notice: For those states with nitrogen application restrictions, including but not limited to, MD, NH, NJ, and VA, do not apply more than 3.5 lbs of this product per 1,000 sq. ft. Dithiopyr: 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S-,S-dimethyl ester . .

EPA Reg. No. 9198-121 EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A, 9198-IN-3S (Underlined letter is first letter used in run code on end of bag.)

KEEP OUT OF REACH OF CHILDREN PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice. If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a

glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person. If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical

ENVIRONMENTAL HAZARDS This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not

Do not contaminate water when disposing of equipment washwaters. It is a violation of Federal Law to use this product in any manner inconsistent with its labeling.

use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. New York Use Restrictions: In New York State this product may only be used by professional

applicators and at no more than 7.0 pounds per 1,000 square feet (equal to 0.5 pounds active ingredient per acre) per year; and is prohibited from use in Nassau and Suffolk Counties. GENERAL INFORMATION

This product is a herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

Turfgrass Safety: This product may be used on seeded, sodded, or sprigged lawns and

ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product.

Cool-Season Grasses Bentgrass, creeping* (Agrostis palustris) Bahiagrass (Paspalum notatum) Bluegrass, Kentucky (*Poa pratensis*) Bermudagrass (*Cynodon dactylon*) Fescue, fine** (Festuca rubra)

Fescue, tall (Festuca arundinacea)

Buffalograss*** (Buchloe dactyloides) Carpetgrass (Axonopus affinis) Ryegrass, perennial (Lolium perenne) Centipedegrass (Eremochloa ophiuroides) Kikuyugrass (Pennisetum clandestinum)

Zoysiagrass (Zoysia japonica)

St. Augustinegrass (Stenotaphrum secundatum)

*Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen',

'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf' and 'Waldina'. Not all varieties of fine fescue have been improved weed control may result. Use of split (half-rate) applications spaced 6-8 weeks apart

Note: It is recommended that use of this product be made on a trial basis initially to determine its F002091 safety under local use conditions and turf management practices before making widespread use. Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

ment until the turfgrass is fully green and has established new roots.

Other Use Precautions: The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller.

For best results, cultural practices that disturb the soil, such as core-, spike-,or hydro- aerification, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices. Apply this product directly to established lawns or ornamental turfs only. DO NOT apply to flowers,

vegetables, shrubs, or trees. and pets off treated areas until dust has settled.

	APPLICATIO	N DIRECTIONS		
TABL		ed Application Use oduct/1,000 sq. ft.)	Rates*	
PROGRAM	1 2		3	
USE (Turfgrass cut)	Preemergence (high)	Preemergence (low)	Preemergence (low)	
REGION		Postemergence (high)	Postemergence (low)	
	i			
North	2.3	2.6	3.5	
North Transition	2.3 2.6	2.6 3.5	3.5 5.2	

*DO NOT apply more than 7 lbs. per application, and no more than 21 lbs. per year. Postemergence control is limited; see "Crabgrass Control" section below and "Precautions" sections above. Regions include areas listed below.

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, NY, & RI. South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX. Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX. **May require split or sequential applications for full control.

Preemergence and Early Postemergence Control: This product provides "preemergence"

control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs. apply directly to water, or to areas where surface water is present, or to intertidal areas below It can also provide "early postemergence" control of crabgrass during the early stages of the mean high water mark. Do not apply when weather conditions favor drift from treated areas. crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turfs. Postemergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the CONTAINER DISPOSAL: Completely empty bag into application equipment. Nonrefillable Read the entire label before using this product. Use only according to label instructions. Not for crabgrass plants in the lawn or turf. So the practical benefit of this product's additional, early postemergence activity is that (compared to strictly preemergence crabgrass products) it can give the user a 2-4 week longer period of time to make applications and still control crabgrass. Application Frequency and Timing: This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall. DO NOT apply more than 7 lbs. per 1,000 sq. ft. per application and no more than 21 lbs.

Spring Applications: For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 above, depending on the user's location, the turfgrass mowing height, and whether the use is considered to be preemergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

Use Program 1 for preemergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control. Use Program 2 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This

Use Program 3 for preemergence control at sites where (a) turfgrass is cut low (e.g, golf year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control at sites where turfgrass is cut low (e.g., golf

Subsequent, sequential pre- and/or postemergence applications should be made where longer Where split fertilizer applications are recommended, the rates in Table 1 may be split across

Dimension is a registered trademark of Dow AgroSciences LLC. two applications made 6-10 weeks apart and prior to crabgrass emergence.

Fall Applications: This product can also be applied in the late summer or early fall (late August 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass bentgrass have been tested. DO NOT apply this product to Colonial Bentgrass (Agrostis tenuis) through the early part of the next spring. The fall application should be followed by an appropriate

following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', and delay mowing again for a few days after the application. When treated lawn or ornamental turf 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'laylo', 'Jamestown', 'Koket', 'Majenta', 'Mary', areas are watered or received significant rainfall within a few days after application of this product. may provide improved weed control.

> Spring Applications: Used as directed for crabgrass control in the spring, this product will also control (at the "Program 3" rates) or suppressed (at the "Program 2" rates) the following weeds

when applied prior to their emergence: barnyardgrass, bluegrass (annual, roughstalk), crowfootgrass, foxtail (yellow & green)*, goosegrass, kikuyugrass, ryegrass (annual & perennial), smutgrass, lespedeza (common), oxalis (buttercup, creeping & yellow woodsorrel), purslane (common), speedwell (corn), spurge (prostrate & spotted). *Also controlled at the "Program 2" rates.

Fall Applications: Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence: bluegrass (annual) - Poa annua, bittercress, chickweed, henbit, geranium (Carolina), parsley-piert,

APPLICATION EQUIPMENT AND INSTRUCTIONS Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more DO NOT use clippings from treated turf for mulching around vegetables or fruit trees. Keep people uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning

	SUGG	ESTED SPREAD	DER SETTIN	GS		
		der settings are r			ration.	
		te your spreader				
50 lbs. covers 21,700 sq. f 50 lbs. covers 19,200 sq.						
50 lbs. covers 14,200 sq. ft						
50 lbs. covers 9,600 sq. f						
			SPREADER SETTINGS			
SPREADER	GROUND SPEED	WIDTH OF COVERAGE	LIGHT RATE 0.16 lbs. ai/A	LOW RATE 0.19 lbs. ai/A	MEDIUM RATE 0.25 lbs. ai/A	HIGH RATE 0.37 lbs. ai/A
Andersons Model 2000/AccPro 2000; (cone 4), Andersons 2000 SR/SR-2000 (cone 4)	3 mph	10 feet	K 1/4	K 1/2	L 1/2	N 1/2
Andersons LCO-1000	3 mph	10 feet	K 1/4	K 1/2	L 1/2	N 1/2
Scotts R-8, R-8A (cone 4)	3 mph	10 feet	J	J 1/4	K 1/2	M 3/4
Lesco 020093 (pattern 0.25)	3 mph	7 feet	D 1/2	D 3/4	Е	F
Lesco 705698 (pattern 0.25)	3 mph	7 feet	11	11 1/2	12	14
Lely Models WTR, WFR, HR, 1250 (PTO at 450 rpm) Pattern II-C	4.5 mph	36 feet	4 3/4	5	5 3/4	7
Vicon (03 Series)	4.5 mph	30 feet	22	25	29	36

STORAGE AND DISPOSAL Do not contaminate water, food, or feed by storage and disposal.

STORAGE: Keep container closed and away from food, feedstuffs, and domestic water supplies. **PESTICIDE DISPOSAL**: Pesticide waste are toxic. Improper disposal of excess pesticide is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

container. Do not reuse or refill this container. Offer for recycling, if available. If not available, then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NOT OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSE-QUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not

program may also be used for early postemergence control at sites where turfgrass is cut high

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Manufactured and Guaranteed by:

The Andersons, Inc. P.O. Box 119 Maumee, OH 43537 www.andersonsplantnutrient.com



The Andersons, Inc. The Andersons, Inc. www.andersonsplantnutrient.com

> 10003593 30000729

010824

Andersons Turf Products 18-0-4

Fertilizer with 0.164% DIMENSION® Turf Herbicide

SEAL AREA NO COPY **SEAL AREA NO COPY**

SEAL AREA NO COPY

SEAL AREA NO COPY

©2024 The Andersons logo is a registered trademark of