Outside of Bag

Acceptable Application Method Noted by an (X)

Only apply this product to established ornamentals.

non-target plants unless they are listed on this label.

Treatment of Ornamental species not listed on this label:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

©2019 The Andersons, Inc. Maumee, OH 43537

10005667

Net Wt. 50 lbs. (22.68 kg)

Do not apply this product directly to bare root of ornamental plants as injury may result. Following transplanting, care must be taken that soil or planting mixes have settled firmly through irrigation, rainfall or packing and that there are no cracks that would allow direct contact of this product to the plant roots or

• Do not apply around ornamental plants that are under stress due to drought, flooding, excessive fertilizer or

soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due

Do not apply under conditions that would affect product distribution. Wind could affect distribution and move product onto sensitive species. Uneven product distribution will result in uneven weed control.
 Do not apply this product on grasses grown for seed.

possible growing conditions. Users who wish to use this product on ornamental species not listed on this label may determine the suitability for use by treating a small number or ornamental plants at a listed rate. Prior to

treatment of larger areas, treated plants should be observed for any symptoms of herbicidal injury, such as foliar damage, reduced vigor or stand reduction, during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious.

When applied as directed under the conditions described on this label, ornamentals listed on back have

shown tolerance in field trials. However, this product has not been tested on all cultivars of each species or under all possible growing conditions prior to large scale use.

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase

LEGAL RIGHTS:THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

 ${\tt CONTRARY\ TO\ LABEL\ DIRECTIONS.\ Use\ contrary\ to\ label\ directions\ is\ not\ permitted.}$

• Do not apply another preemergence herbicide within 4 weeks of application of this product.

Acceptable Application Method Noted by an (X)

Ornamentals Noted by an (X)

A cceptable Application Method Noted by an (X)

15.50"



Professional Andersons Turf Products

Dimension ® 0.25G

• For use on Turf, Landscape Ornamentals, and Nursery Ornamentals • On DG Pro® Carrier

areas, established lawns and ornamental turf (including golf course fairways, roughs and tee boxes and sports turf) and container and field grown nursery ornamentals. New York Use Restrictions: In New York State this product may only be used by professional applicators and at DO NOT use clippings from treated turf for mulching around vegetables or fruit trees. no more than 4.6 pounds per 1000 square feet (equal to 0.5 pound active ingredient per acre); and is prohibited from use in Nassau and Suffolk Counties. ACTIVE INGREDIENTS: OTHER INGREDIENTS . .

CAS No. 97886-45-8 *This product contains 0.125 pounds of active ingredient, dithiopyr, per 50 pound bag. EPA Est. No. 9198-OH-1<u>M</u>, 9198-OH-2<u>B</u>, 9198-AL-001<u>A</u> PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS KEEP OUT OF REACH OF CHILDREN

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some

If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison center or doctor for treatment advice.							
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulanc then give artificial respiration, preferably mouth-to-mouth if possible. Call poison control center or doctor for treatment advice.							
If swallowed	Call a poison control center or doctor for treatment advice. Have person si a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.							
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment adv							

going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

This product is toxic to fish and highly toxic to other aquatic organisms including ovsters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters. Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forest, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, tification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protection equipment (PPE), restricted-entry intervals. The requirements in this box only apply to uses of this product, that are covered by the Worker Protection Standard. Do n otherwise specified in the directions for use associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: 1. Coveralls over short sleeve shirt and short pants, 2. Chemical-resistant footwear, shoes plus socks, 3. Chemical-resistant gloves made of any waterproof material, and gloves, 4. Chemical-resistant headgear for overhead exposure. When the Restricted-Entry Interval for a crop is 7 days or longer, you must notify workers of the application by warning them orally and by posting warning signs at entrances to treated area.

rates listed in Table 2 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-long control. Non-Agricultural Use Requirements The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, nurseries, or greenhouses. Do not enter or allow others to enter the treated area (except those involved in the watering-in) until the watering-in is completed and the area has dried.

t is a violation of Federal Law to use this product in any manner inconsistent with its labeling. Read the entire label before using this product. Use only according to label instructions.

STORAGE: Store this product only in its original container in a dry, cool, secured storage area. Do not contaminate water, food, or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. eprocessed should be disposed of in a landfill approved for pesticide disposal in accordance with applicable CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke. SPILL: In case of spill, sweep up material and dispose of material according to "Pesticide Disposal" directions

for research purposes. This product is a herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in product will not harm nearby established ornamentals when used according to label directions.

TURFGRASS SAFETY is product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product: Warm Season Grasses

Bentgrass, creeping* Fescue, tall Bahiagrass Carpetgrass St. Augustinegrass Bluegrass, Kentucky Ryegrass, perennial Bermudagrass Centipedegrass Zoysiagrass Fescue, fine** Buffalograss*** Kikuyugrass *Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. DO NOT apply this product to Colonial bentgrass (Agrostis tenuis) varieties. **Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf', 'How this work of the second of the s

and 'Waldina'. Not all varieties of fine fescue have been tested. ***DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots. user evaluate the response of turfgrass varieties to this product on a limited area, at use rates, prior to initiating

RESEEDING, OVERSEEDING, OR SPRIGGING Reseeding, overseeding, or sprigging of treated areas with this product should be delayed until 12 weeks from the time of application. Reseeding, overseeding or sprigging before 12 weeks after application may prevent establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil equipment (slit seeders) designed to give good seed to soil contact.

APPLICATION DIRECTIONS CONTROL OF CRABGRASS

when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs. It can also provide "early postemergence" control of crabgrass. Postemergence crabgrass control will be product's early postemergence activity is it can give the user a 2-4 week longer period of time to make

nis product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall. DO NOT apply more than 4.6 lbs. per 1000 sq. ft. per application, and no more than 13.8 lbs. per 1000 sq. ft. per

For single applications made in the spring or early summer, this product should be applied at the appropriate rate listed in Table 1 below. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region. Use the higher rates where weed infestations are dense. Also use the higher rates if applications are made early and longer residual control is desired. Subsequent, sequential pre- and/or postemergence applications should be made where longer periods of Where split fertilizer applications are listed, the rates in Table 1 may be split across two application made 6-10

	Weeds Controlled	Application Rate	WA	
	ass: (Large, Smooth, Sou Foxtail (Giant, Green, Ye	108-160 lbs./acre 2.4-3.6 lbs./1,000 sq ft 0.27-0.40 lbs. ai/acre	Mar fit f mal PAF	
Barnyardgrass Chickweed Clover, hop Crabgrass, tropical Crowfootgrass Dichondra, Carolina Lespedeza, common Kikuyugrass Mary's grass Marestail Medic, black	Mulberry weed Mustard Oats, wild Oxalis Buttercup Creeping Woodsorrel Yellow Woodsorrel Pineappleweed Ryegrass, annual Ryegrass, perennial Sandbur	Smutgrass Speedwell, corn Spurge Garden Prostrate Spotted	160-200 lbs./acre 3.6-4.6 lbs./1,000 sq. ft. 0.40-0.50 lbs. ai/acre	LIAI pric DISC COI LEG THII VAF Visi © 20 P.O.
Bluegrass, annual Bittercress Carpetweed Chickweed Geranium Carolina Goosegrass	Henbit Knotweed, prostrate Parsley-piert Pigweed, redroot Purslane, common Rocket, London Shepherdspurse		200 lbs./acre 4.6 lbs./1,000 sq. ft. 0.50 lbs. ai/acre	1(30
he higher rates where w		'		N 6

Table 2: Application Use Rates for Fall Applications (lbs. of product/1,000 square feet)

*DO NOT apply more than 4.6 lbs. per application, and no more than 13.8 lbs. per year. Postemergence control is Tips For Improved Control For best results, apply this product within a few days after mowing and delay mowing again for a few days after When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result. Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control. APPLICATION EQUIPMENT AND INSTRUCTIONS Apply this product with suitable spreaders such as drop or rotary-type designed to apply granular herbicides.

Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate listed above.

Apply this product uniformly over the lawn or ornamental turf area. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform ORNAMENTALS (Landscape, Field Grown and Container Grown) This specialty herbicide provides preemergence control of listed annual grass and broadleaf weeds in areas planted with tolerant ornamental plants listed on this label. It is intended for use on plants grown for aesthetic purposes in landscaped areas or in container or field grown production nurseries. When applied as directed, the

product may be applied as a single application or as a split application. Refer to the following table for product DO NOT apply more than 200 lbs. per acre per application and no more than 3 applications (600 lbs.) per acre In New York State, this product may only be used by Professional Applicators at no more than 200 lbs. per acre (4.6 pounds per 1,000 sq. ft./ 0.5 lbs. active ingredient per acre) per year; and is prohibited from use in Nassau Pounds of Product Pounds of Product Ounces of Product per Acre per 1,000 square feet per 100 Square Feet

200 4.6 APPLICATION TIMING AND RECOMMENDATIONS


This product is a preemergence herbicide that will not control established weeds. Make applications prior to weed seed germination or to bare ground. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Prior to making an application, control existing vegetation by hand weeding, cultivation or with a postemergence herbicide. Once an application of this product has been made, do not disturb the soil surface as the herbicide barrier will be broken.

Pigweed, redroot Pineappleweed

Oxalis, buttercup Woodsorrel, yellow

Foxtail, yellow Bittergrass Carpetweed Geranium, Carolina Purslane, common Crabgrass, Iarge Ryegrass (annual & perennial) Knotweed prostrate Shepherdspurse Crabgrass, smooth Sandbur Lespedeza, common Speedwell, corn Crabgrass, southern Smutgrass Medic, black Spurge, prostrate
Mulberry weed Spurge, spotted
Mustard Woodsorrel, creeping





SEAL AREA NO COPY

PANTONE 725 C

PANTONE 130 C

	Sugges	ted Spreader	Settings
These suggested spreader settings are n	ot intended to	replace calibr	ation. Calibrate your spreader before applying product.
50 lbs treats 20,8	00 sq. ft. (0.4	8 acre) at the l	LOW RATE (2.4 lbs/1,000 sq .ft.)
50 lbs treats 14,700	sq. ft. (0.34 a	acre) at the ME	DIUM RATE (3.4 lbs/1,000 sq. ft.)
50 lbs treats 13,80	00 sq. ft. (0.32	2 acre) at the H	IIGH RATE (3.6 lbs/1,000 sq.f t.)
50 lbs treats 10,80	0 sq. ft. (0.25	acre) at the HI	EAVY RATE (4.6 lbs/1,000 sq. ft.)
			SDDFADED SETTINGS

			SPREADER SETTINGS							
SPREADER	GROUND SPREADER	WIDTH OF COVERAGE	LOW RATE 2.4 lbs/1,000 sq. ft.	MEDIUM RATE 3.4 lbs/1,000 sq. ft.	HIGH RATE 3.6 lbs/1,000 sq. ft.	HEAVY RATE 4.6 lbs/1,000 sq. ft.				
Andersons Model 2000/ AccuPro 2000; (cone 6) Andersons 2000 SR/ SR-2000 (cone 6) Andersons LCO-1000	3 mph	11 feet	К	L 1/4	L 3/4	M 1/2				
Scotts R-8, R-8A (cone 6)	3 mph	11 feet	J	K	K 1/4	L 1/2				
LESCO 020093 (0.25)	3 mph	7 feet	D	Е	E 1/4	F				
LESCO 705698 (0.25)	3 mph	7 feet	10	12	12 1/2	14				
Lely Models WTR, WFR, HR, 125- (PTO at 450 rpm)	4.5 mph	34 feet	4 1/4 IIIC	5 1/4 IIIC	5 1/2 IIIC	6 IIIC				
Vicon (03 Series)	4.5 mph	30 feet	18	23	26	29				




(Abelia x grandiflora) abyssinian red banana (Ensete ventricosum)	maureli	Х	Х	×	Х	(Asparagus spp.) fescue (Festuca glauca)			Х		Х	(Narcissus spp.) oak, laurel			X		+
acacia redolens (Acaia redolens)		Х	Х	Х	Х	fetterbush (Leucothoe fontanesiana)	rainbow		Х		Х	(Quercus laurifolia) oak, pin		 	X		+
Ajuga carpet bugle	bronze bronze beauty		×		X	ficus (Ficus retusa)	nitidia		Х		Х	(Quercus palustris)					\bot
(Ajuga genevensis)	bronze beauty					fir fraser			Х		Х	oak, red (Quercus rubra)		<u> </u>	X		\perp
almond, flowering (Prunus gladulosa)			Х		Х	(Abies fraseri) Forsythia						oak, southern (Quercus virginiana)			X		4
apple† (Malus pumila)			X		Х	(Forsythia x 'Arnold Dwarf') (Forsythia viridissima) (Forsythia xintermedia)	arnold dwarf bronxensis dwarf		X X		X X	oak, willow (Quercus phellos)			X		\perp
	nigra pyramidalis		X X X		X X X	(Forsythia x 'Meadowlark')	lynwood gold meadowlark		X X		X X	oleander (Nerium oleander)	hardy red petite pink		X		
	smaragh techny woodwardii		X X		X X	(Forsythia x intermedia)	spring glory	X	×	×	x	osteospermum	sister agnes whirligig	 	X		\dagger
	aurea nana	Х	Х	Х	Х	(Forsythia suspensa)	weeping		Х		Х	(Osteospermum fruticosum) pachysandra	<u> </u>	 	X		+
(Thuja orientalis) ash, green			X	-	Х	fountain grass, purple (Pennisetum setaceum)	rubrum	X	Х	Х	X	(Pachysandra terminalist) palm, King	<u> </u>	 	X		+
(Fraxinus pennsylvanica) ash, autumn purple	autumn purple		X	-	X	fuchsia (Fuchsia spp.)			Х		X	(Archontophoenix cuninghamiana)					
(Fraxinus americana)	dwarf queen		X		X	galium (Galium ordoratum)			Х		Х	palm, queen (Syagrus romanzoffianum)		Х	Х	×	T
(Callistephus chinensis)	-					gardenia (Gardenia jasminoides)	mystery radicans		X X		X	pampas grass (Cortaderia selloana)			Х		\dagger
azalea (Rhododendron spp.)	brilliant buccaneer carror	X X X			X X X		white gem	X	×	X X	X	pansy		 	Х		\dagger
	chimes (Belgian)	X			X	Garlic, variegated society (Thulbaghia violacea)	variegata	_ ^		^		(Viola x wittrockiana) paper flower	Barbara Karst	X	X	X	十
	Elsie Lee exbury	X X			X X	geranium (Pelargonium x hortorum)			Х		Х	(Bougainvillea glabra) peach†	<u> </u>	 	X		+
	fashion hardijzer	X			X	gum (Eucalyptus citriodora)			X		Х	(Prunus persica) periwinkle, dwarf		 	X		+
	beauty hersey red higasa	X X			X X	hawthorn (Crataegus spp.)	cockspur white crimson cloud		X X		X X	(Vinca minor)			ļ		\bot
	hinocrimson Holland	X			X		enchantress Jack Evans		X X		X X	petunia (Petunia x hybrida)	picoti	ļ	×		\downarrow
	(hybrid) Marion Lee	×			X		Washington white		Х		X	photinia, red tip (Photinia x fraseri)			X		\perp
	northern lights orange cup orchid lights	×			X X	hawthorn, Indian (Rhaphiolepsis indica)	ballerina		Х		Х	pieris (Pieris taiwanensis)			X		
	snow southern	X			X	heather, twisted (Erica cinerea)	Mediterranean pink		Х		Х	pieris, Japanese (Pieris japonica)	mountain fire		X		
azalea, flame	charm	Х	X		X	hemlock, Canada (Tsuga canadensis)			Х		Х	pine, Afghan (Pinus eldarica)		Х	Х	×	T
(Rhododendrom calendulaceum)					^	hisbiscus (Hibiscus spp.)	blue bird brilliant		X X		X X	pine, Austrian black (pinus nigra)		Х	Х	×	\dagger
azalea, kirishima (Rhododendron obtusum)			Х		Х		hula girl		x		x	pine, Japanese black		+	Х		十
bamboo heavenly	compacta	<u> </u>	X		Х	holly (llex x 'Nellie R. Stevens')	Nellie R. Stevens	×	×	×	x	(Pinus Thunbergiia) pine, loblolly			X		+
	nana		X		X	(llex x attenuata)	fosteri Savannah		X X		X X	(Pinus taeda) pine, longleaf			×		+
	dwarf pygmy green		X X		X X	holly, blue (llex x meserveae)	blue bird brilliant		X		X	(Pinus palustris)		<u> </u>	X	ļ	4
	kobold pygmy red rose glow		X X X		X X X		brilliant hula girl		Х		Х	pine, mugo or Swiss Mt. (Pinus mugo)			ļ		\downarrow
barberry, purple	rose glow atropurpurea		X		X	holly, cassine (llex cassine)			X X		X X	pine, Scotch (Pinus sylvestris)			×		\perp
(Berberis thunbergii var atropurpurea)						holly, Chinese (Ilex cornuta)	burfordii needlepoint		X X		X X	pine, slash (Pinus elliottii)			×		
basket flower (Gaillardia grandiflora)			Х		Х	holly, Japanese (llex crenata)	compacta hellerie		X		X	pine, Virginia (Pinys virginiana)			X		T
beach grass (Ammophila breviligulata)			Х		Х		Japanese northern beauty		X		X	pine, white (Pinus strobus)		X	×	×	+
bearberry (common) (Arctostaphylos uva-ursi)	Massachusetts		Х	İ	Х	holly, yaupon	steeds	×	X	Х	X	pineapple guava†			×		+
bee balm			Х		Х	(llex vomitoria) honeysuckle						(Feijoa sellowiana) pittosporum, tobira	1		 		+
(Monarda didyma) begonia			Х		Х	(Lonicera xylosteum) (Lonicera japonica)	Claveyi dwarf halliana		X X		X X	(Pittosporum tobira)	Wheeler's dwarf	Х	×	×	\rfloor
(Begonia spp.) birch river		<u> </u>	X	-	X	(Lonicera tatarica)	Canadian white Zabelli		X X		x x	plumbago, cape (Plumbago auriculata)	royal cape	×	×	×	Γ
(Betula nigra) Blackeyed Susan	goldstrum		X	-	×	hosta (Hosta sieboldii)			Х		X	potentilla (Potentilla fruticosa)	abbotswood		×		+
(Rudbeckia hirta)	gordattutti	<u> </u>				(Hosta lancifoliai)	albo marginata		Х	×-	Х	(Potentilla nepalensis)		 	X		+
blanket flower (Gaillardia spp.) blue fescue			X X		X X	ice plant (Carpobrotus edulis)		X	Х	Х	Х	privet (Ligustrum x vicaryii) (Ligustrum japonicum)	golden vicary regal		××		
(Festuca ovina)	la la constantina de		ļ			ice plant (Drosanthemum floribundum)		X	Х	Х	Х	(gastram japonicum)	texanum yellow tipped	Х	X X	×	
blueberry† (<i>Vaccinium spp.)</i>	bluecrop blue jay		X		X	impatiens (Impatiens spp.)			Х		Х	privet, glossy (Ligustrum lucidum)	3		Х		\dagger
	Jersey north blue northland		X X X		X X X	iris (iris spp)	dwarf blue wedgewood		X X		X X	Pyracantha or firethorn		<u> </u>			\dagger
bottlebrush	Little John	Х	X	Х	X	ivy, English	Bulgaria		Х		Х	(Pyricantha x 'Gnome') (Pyricantha coccinea)	gnome lalandei		X X X		
(Callistemon citrinus) bougainvillea	James Walker	<u> </u>	X		Х	(Hedera helix) jasmine, Asian	thorndale		X		X	(Pyricantha koidzumii) queen palm	victory	X	X	X	+
(Bougainvillea spectabilis) boxwood	Scarlet O'Hara japonica	X	X	X	X	(Trachelospermum asiaticum) jasmine, star			X		X	(Arecastrum rammanzoffianum)					\perp
(Buxus microphylla)		X	X	X	X	(Trachelospermum jasminoides)			Α			quince, Japanese † (Chaenomeles japonica)			X		
boxwood, green beauty (Buxus microphylla japonica)	green beauty			<u> </u>		juniper (Juniperus spp.)	Arcadia Armstrong	İ	X		X	Rhododendron (Rhododendron spp)	album Cunningham		X		Т
boxwood, welleri (Buxus sempervirens)	winter gem	X	X	×	Х	(Jumperus Spp.)	bar harbor blue chip		X		X X		white PJM		×		
broom (Cytisus scoparius)	moonlight		×		X		blue Pacific blue rug		X X		X X		purple gem silvery pink		X X		
(Genista pilosa) camellia	Vancouver gold	<u> </u>	X		X		blue star broadmoor		X		X	rhododendron, Carolina (Rhododendron			X		
(Camellia japonica)	debutante mathotiana	×	X X	×	X X		buffalo calgary carpet		X X X		X X	carolinianum) rhododendron, catawba		 	X		+
(Camellia sasanqua)	supreme chansonette	×	X X	×	X X		emerald sea emerald spreader		X		×	(Rhododendron catawbiense)			_ ^		
	setsukgekka snow white	<u> </u>	X		X		endora com pacta		Х		X	rhododendron, rhodie max- rosebay			Х		\dagger
(Iberis sempervirens) carex, variegated		<u> </u>	X		X		fruitlandi green		X X		X X	(Rhododendron maximum)		 			\downarrow
(Carex spp.)			-				gold tip hetzi		X X		X X	ribbon grass (Phalaris arundinacea)		ļ	X		\perp
	bronze bronze beauty		X X		X X		hughes Manhatten blue parsoni		X X		x x	rockcress (Arabis caucaisca)	snowcap		×		\perp
cedar, red			Х		Х		pfitzeriana plumosa		X		X	rose† (Rosa banksiae)	luta		X		
(Juniperus virginiana) Celosia		<u> </u>	X		Х		Prince of Wales procumbens		X X		X X	rose, knockout shrub (Rosa spp. hybrid)	knockout	×	Х	×	Т
(Celosia spp.) Centaura		<u> </u>	X		Х		dwarf San Jose sargent blue		X X		X X	rosemary† (Andromeda officinalis)			Х		T
	scarlet		×		×		sargent green scandia		X		X X	rosemary, bog	nana		Х		\dagger
coleus	red kewpie		X		Х		scopulorum moonglow		Х		x	(Andromeda polifolia) salvia (Salvia farinacca)	rhea		×		+
(Coleus blumei)		<u> </u>	X	-	X		spartan tamariscifolia		X X		X X	(Salvia farinacea) sedum	dragon blood		×		+
(Aquilegia spp.)			X	-	×		weberi Youngstown Yukon belle		X X		X X	(Sedum spurium)	red red carpet		×		
coneflower, purple (Echinacea purpurea)	magnus purple	Х	Х	Х	Х	juniper	gold coast	×	Х	X	Х	smoketree, royal purple	yellow royal purple		X		+
copper leaf (Acalypha wilkesiana)	-		X		Х	(Juniperus chinensis)	torulosa-Holly	×	Х	×	×	(Cotinus coggygria) snapdragon		+	×		+
coreopsis (Coreopsis spp.)	moonbeam		×		Х		wood twisted					(Antirrhinum spp.)	sterile	×	X	×	+
corn flower (Centaurea spp.)		L_	Х	<u> </u>	Х	juniper (Juniperus procumbnes nana)	prostrate Japanese	×	×	×	X	snowball, common (Viburmum opulus)	Storing	_ ^		_ ^	4
cotoneaster (Cotoneaster apiculatus)			Х		Х	juniper	garden tamarix	×	X	X	X	sourwood (Oxydendrum arboreum)		<u> </u>	X		\perp
coyotebrush (Baccharis pilularis)			×		Х	(Juniperus sabina 'tameriscifolia')					<u> </u>	spiraea (Astilbe X arendsii)	fanall		×		
cycad			×		Х	juniper (Juniperus scopulorum)	scopulorum	×	Х	×	х	spiraea (Spiraea spp.)	Anthony Water red		Х		
(Cycas revoluta) cypress, bald			×	 	Х	king palm	skyrocket	X	X	X	X		dolchica froebeli pink		X X		
(Taxodium distichum)	glauca		X		X	(Archontophoenix cunninghamiana)	<u>L</u> _	L_	_		_		goldenflame red		×		
(Cupressus sempervirens)	gracilis		X	-	×	laurel, Carolina cherry (Prunus caroliniana)	bright n' tight	×	Х	Х	Х		snowmound white Van Houtte		×		
(chamaecyparis obtusa)	g. acm3					laurel, moutain (Kalmia latifolia)		İ	Х		Х	spiraea garland	white		X		+
cypress, leyland (Cupressocyparis leylandii) hybrid			X		X	leucothoe		İ	×		Х	spiraea, garland (Spiraea x arguta)		<u> </u>	ļ	<u> </u>	+
daffodil	King Alfred		×		Х	(Leucothoe fontanesiana) ligustrum, Japanese			X		Х	spruce, Black Hills (Picea glauca var densata)	1.	<u> </u>	X		\downarrow
	Aztec gold		X		X	(Ligustrum japonicum)	steamline		X		Х	spruce, Colorado blue (Picea pungens)	glauca	×	×	×	\perp
	bright yellow (hybrid)		×		X	(Agapanthus spp.) Lily of the Nile	albus	<u> </u>	X		X	spruce, Norway (Picea abies)			Х		
	single gold (evergreen) Wilson's		×		X X	(Agapanthus africanus)	Peter Pan	<u> </u>	Х		Х	spruce, white (Picea glauca)	conica		×		T
dianthus (owest MC)	yellow				× ×	lilyturf (Liriope muscari)	evergreen giant lilac beauty	×	X X	¥	X X	sweetflag grass (Acorus calimus)			X	İ	†
dianthus (sweet William) (Dianthus spp.)	-		X		^		majestic monroe white	×	X X	×	X X X	sweetgum			X		+
delphinium (Delphinium spp.)	magic fountain		×		Х	<u></u>	silvery sunproof variegata	<u></u>	X X		X X	(Liquidambar styraciflua) sycamore, American	 	 	×		+
dogwood (Cornus florida)		<u> </u>	Х	<u> </u>	Х	lilyturf, creeping (Liriope spicata)			Х		Х	(Platanus occidentalis) tea tree, New Zealand	ruby glow	X	×	×	+
+	flavarimea		×		Х	maple, amur	emerald elf	×	Х	Х	Х	(Leptospermum scoparium)	, 9,0**	<u></u>	ļ		4
Douglas fir			X		Х	(acer ginnala) maple, Japanese			X		Х	tree fern (tiki fern) (Asparagus virgatus)			X		\perp
(Pseudotsuga menziesii) dusty miller		X	X	Х	Х	(Acer palmatum) maple, Norway			X		Х	trumpet flower or Carolina jessamine			×		
	drake		×	-	Х	(Acer platanoides)	red sunset	×	×	X	X	(Gelsemium sempervirens)	apeldoorn	 	×		+
(Ulmus parvifolia) eulaliagrass	maiden grass	×	X	X	X	(Acer rubrum)	. ca sunset			^		(Tulip spp) tufted hairgrass		+	×		+
(Miscanthus sinensis)		_ ^	_ ^	_ ^	_ ^	maple silver (Acer saccharinum)			Х		Х	(Deschampsia caespitosa)	 				4
euonymus (Euonymus japonicus)	Argenteo- variegata		×		×	maple sugar† (Acer saccharum)	L	L	×		Х	verbena, shrub (Lantana sellowiana)		<u> </u>	X	ļ	\perp
(Euonymus fortunei)	Auereo- marginata		×		×	marigold (Tagetes patula)	honeycomb variegata		X		X X	Verbena, St. Paul's (Verbena peruviana)	St. Paul		×	L	
	colorata emerald gaiety emerald n' gold		X X		X X	(.agetes patuid)	wheeleris dwarf		X X		x x	viburnum (Viburnum spp.)	American cran- berry bush		×		†
	emerald n' gold gold edge gold princess		X X		X X X	mock orange†	golden		X		X	Spy)	arrowood European cran-		×		
	gold princess Manhattan	×	×	×	x x	(Philadelphus spp)	snowflake double white	ļ	Х		X		berry bush linden		×		
I	silver king	"	×		x	mondo grass (Ophiopogon japonicus)	<u> </u>		Х		Х	<u></u>	Mohican wright		×	<u>L</u>	
I	tricolor vegetus	<u></u>	x x	<u>L</u>	x x	moss rose (Acer rubrum)	sunnyside		Х		Х	vinca (periwinkle) (Vinca minor)			×		\top
	viridis	×	×	Х	X	mountainash			Х		Х	windmill palm		\vdash	×		+
euryops, green leaved			×		Х	(Sorbus aucuparia) myrtle, crape	faurei		X		X	(Trachycarpus fortunei) xylosma	 		×		+
euryops, green leaved (Euryops pectinatus) fan palm, European			×	1	Х	(Lagerstroemia indica)	langer muskogee		X X		X X	(Xylosma congestum) yarrow	-		X		+
euryops, green leaved (Euryops pectinatus)		ı		1	ı l	1	standard pink	I	X		X	(Achillea spp.)	1	I	1 ^	I	
euryops, green leaved (Euryops pectinatus) fan palm, European (Chamaerops humilis)			<u> </u>			myrtle, wax			Х		X		al 1	+		-	+
euryops, green leaved (Euryops pectinatus) fan palm, European (Chamaerops humilis) fan palm, Mexican			I	1		myrtle, wax (Myrica californica)			Х		Х	yaupon (llex vomitoria)	dwarf		Х		1

SEAL AREA NO COPY

1620 Tech Avenue, Mississauga, ON L4W 5P4 Canada (905) 219-1600 Please check carefully before using. Although every effort is made to produce the finest possible product, errors do occur. We do not assume any liability beyond the replacement of our product. SEQ 100% 75% 50% 25% 10% 3% 2% 1% Color Client Name: Direct Plastics Docket: 936270A Item Part #: 01911FG 05 / 10 / 2018 Bar Width Adi.: - 0.005" 031865315381 Line Screen: 65 Print Method: Flexo Project:



PLEASE CHECK CAREFULLY!



SEAL AREA NO COPY

AND009371

.375 Seal

SEAL AREA NO COPY