TURF AND ORNAMENTAL SYSTEMIC FUNGICIDE

For the prevention and control of turf diseases and the diseases of annual and perennial flowers, bedding plants, foliage plants, ground covers, plus deciduous and evergreen trees and shrubs.

ACTIVE INGREDIENT Thiophanate-methyl OTHER INGREDIENTS

Keep Out Of Reach Of Children **CAUTION**

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or when going for treatment. For 24-Hour Emergency Assistance Call Chemtrec 1-800-424-9300.

EPA Reg. No. 1001-70



Cleary Chemicals LLC 178 Ridge Road, Suite A, Dayton, New Jersey 08810-1501 (732) 329-8399 (800) 524-1662

NET CONTENTS: 30 lb. (13.6 kg.)

FPO 0416410010030 3 of 9 Barcode Type



CLETP3B

SEAL AREA NO COPY

TURF AND ORNAMENTAL SYSTEMIC FUNGICIDE

Keep Out Of Reach Of Children IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the Apply 2 treatments at 21 day intervals in late fall to early winter, with the last application made just prior to first permanen snow cover. Rotations with PCNB can be utilized. EPA Reg. No. 1001-70 EPA Est. No 9198-OH-1<u>M</u>, 9198-OH-2<u>B</u>, 9198-AL-001<u>A</u> To Ridge Road, Sulte A, Dayton, New Jersey 08810-1501 (732) 329-8399 (800) 524-1662 (Underlined letter is first letter used in run code on the end of bag) with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Loaders, applicators and other handlers must wear: 6-9 Apply when disease first appears at 14 day intervals. Chemical resistant gloves are required for applicators using hand held equipment Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Apply middle-late Fall to early Winter before turf has stopped all growth activity or in spring during periods of thaw, or when disease symptoms first appear. Continue at 14 day intervals throughout anticipated disease period. Best results are achieved when first application is made before turf is completely dormant.

Apply when disease first appears and make two applications at 14 day intervals. For prevention, apply in spring and fall. pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove clothing/PPE immediately after handling this product. As soon as possible, wash thoroughly and 6-9 For prevention, apply 3 applications starting late April or early May using 21 day intervals. Rotations may be used as part of the three application program. For suppression, apply at 14 day intervals when disease first appears.
6-9 For prevention, apply in early June or based upon local Extension Service recommendations. Apply at 14 day intervals as needed. Rotations and/or tank mix combinations may be used for season long disease prevention. Do not apply directly to water, or areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any other requirements *Observe the maximum individual application rates and maximum seasonal application limits in Table 2. AGRICULTURAL USE REQUIREMENTS Table 4: Guide to 3336 DG Lite Recommended Spreader Settings* Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific These settings are presented as a guide for use. Age, condition of spreader and operator speed will cause variation in applied rates of material. Spreader calibration at the beginning of each application is strongly recommended. statements and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval (REI). The requirements in this box apply to uses of this

| Spreader | Speed (MPH) | Swath (FT) | Cone Set | Rate of 3336 DG Lite lbs/1,000 sq ft | 3 | 6 | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 | | 9 |

COMMERCIAL HORTICULTURAL APPLICATIONS

Annual and Perennial Flowers, Bedding Plants, Foliage Plants, Ground Covers, plus Deciduous and Evergreen Trees and Shrubs

3336 DG Lite is a broad spectrum systemic fungicide which controls a variety of foliar, stem, and root diseases on a wide range of commercially important plants. 3336 DG Lite may be used for disease control and prevention programs on annual and perennial flowers, bedding plants, foliage plants, ground covers, and deciduous and evergreen trees and shrubs found in Nursery, Greenhouse, Interiorscape, and Landscape sites. Follow guidelines provided under general information for use of this product.

activated with irrigation or natural rainfall within 24 hours for best results. If irrigated, application should be watered in by applicator with 0.1-0.25 inches of water to begin activation. For media incorporation, mix at a rate of 2-3 lb/cu yd before seeding or transplanting of desired crop. Media incorporated with 3336 DG Lite may be stockpiled for later use without activation until needed.

or implied warranty. To the extent consistent with applicable law, neither CLEARY CHEMICAL nor seller shall be liable for consequential, special or indirect damages resulting from the use or handling of this product including, but not limited to, loss of profits, business reputation, or customers, labor costs, or other expenses incurred in planting or harvesting. CLEARY CHEMICAL and seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized

over the top application after seeding or transplanting

to propagation beds, containers, pots, or nursery or landscape beds. For curative control, apply after

seeding or transplanting when disease first appears. Repeat every 21-28 days for adequate crop protection. Note: 3336 DG Lite does not control Pythium or Phytophthora. Rotations with metalaxyl, mefenoxam etridiazole, propamocarb, mono and dipotassium salts

Note: The "Directions For Use" of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on Swedish Ivy (Plectranthus australis), Boston Ferr (Nephrolepis exhalta), and Easter

Nursery, Greenhouse, Landscape & Interiorscape

For control of soil borne diseases effecting Bulbs, Corms, Rhizomes, Roots, Crowns

and Stems caused by: Botrytis, Cylindrocladium, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia,

DG Lite logo is a registered trademark of The Andersons, Inc. 3336 is a registered trademark of Cleary Chemicals, LLC.

LIMITED WARRANTY AND DISCLAIMER

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours. Personal Protective Equipment (PPE) required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, · Coveralls over long-sleeved shirt and long pants · Chemical-resistant gloves made of any waterproof materia · Chemical-resistant headgear for over head exposur

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this

STORAGE AND DISPOSAL Do not contaminate water, food, or feed by storage or disposal. Pesticide Storage: Store in the original container in a dry, temperature controlled area. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling, contain/re-capture spillage and use in accordance with instruction below or dispose of in accordance with the Pesticide Disposal Instructions listed below.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be used according to the label Container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Shake empty product container to remove all possible material. When bags are empty, they may be disposed of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

COMMERCIAL TURF AND ORNAMENTAL

Apply 3336 DG Lite with ground based, hand operated (push) or mechanical (power) equipment sufficient to accurately deliver this product to the application site. Do not apply by hand or with hand held rotary spreader (e.g. bellygrinder). Application equipment should be properly calibrated prior to use. Application equipment should be thoroughly cleaned after use. If a rate range is given, use the meowner use and can only be used by individuals/firms licensed or registered by the state to apply omamental or turf pest control products. No claim of compatibility with other pesticides is implied. If, after using 3336 DG Lite as recommended, the treatment is not effective, a tolerant strain of fungi may be present. Consult your local Cleary Chemical representative, your State Agricultural Experiment Station, or your State Cooperative Extension Service for proper disease identification and advice on the prompt use of some other suitable fungicide or disease control strategy. As long as recommended precautions are followed, 3336 DG Lite can remain useful for disease control. COMMERCIAL TURF APPLICATIONS For use on all fine turf applications such as commercial, residential and public (such as home lawns, parks, athletic fields, schools, and day care centers), and golf courses (greens, tees, fairways and aprons) of cool and warm season grasses such as Bentgrass, Bluegrass, Bermudagrass, Fescue, Ryegrass, St. Augustinegrass, Zoysiagrass, or their mixtures. 3336 DG Lite is not phytotoxic to any of the above mentioned grasses when used in accordance with the label. 3336 DG Lite is to be used for the prevention and control of the diseases mentioned

under conditions of severe disease pressure. Do not graze animals on treated turf. Do not feed clippings to livestock or poultry. Not for use on turf being grown for sale or other commercial use as sod. Table 1: Maximum Individual Application Rates and Minimum Re-Treatment Intervals

Use Site	Maximum application rate of 33	m application rate of 3336 DG Lite Re-Treatmen		Comments
Residential or Public Areas	131 Lbs/Acre (3 lbs / 1,000 :	sq.ft.) 14	days	
Golf CourseTees, Greens, Aprons	392 Lbs/Acre (9 lbs / 1,000	sq.ft.) 14	days	
Golf Course Fairways – except Florida	262 lbs/Acre (6 lbs / 1,000 :	sq.ft.) 14	days	Excludes Florida
Golf Course Fairways - Florida Only	131 lbs/Acre (3 lbs / 1,000 :	sq.ft.) 14	days	Florida Only
ble 2: Maximum Seasonal Applica	Maximum Pounds 3336 DG	following amounts of produ Pounds 3336 DG Lite	' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' 	
			' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' 	
ble 2: Maximum Seasonal Applica Use Site Residential or Public Areas			Comments 4 Applicatio	
Use Site Residential or Public Areas	Maximum Pounds 3336 DG Lite per Acre per Season 523 lbs.	Pounds 3336 DG Lite per 1,000 sq.ft. 12 lbs.	Comments	
Use Site Residential or Public Areas Golf Course Tees, Greens, Aprons	Maximum Pounds 3336 DG Lite per Acre per Season 523 lbs. 1048 lbs.	Pounds 3336 DG Lite per 1,000 sq.ft. 12 lbs. 24 lbs.	Comments 4 Applicatio	ns per year
Use Site Residential or Public Areas	Maximum Pounds 3336 DG Lite per Acre per Season 523 lbs.	Pounds 3336 DG Lite per 1,000 sq.ft. 12 lbs.	Comments	ns per year

For prevention in historic areas of disease pressure, apply twice at 14 day intervals when soil temperature reaches 60°F. For curative control, apply when disease first appears and continue at 14 day intervals. Rotations with chlorothalonil or triadimefon can be utilized.

The Andersons and DG Lite logo are registered trademarks of The Andersons, Inc. 3336 is a registered trademark of Cleary Chemicals, LLC.

FPO 0416410010030 3 of 9 Barcode Type

NET CONTENTS: 30 lb. (13.6 kg.)



SEAL AREA NO COPY