



- Selective Preemergent Herbicide for the Control of Annual Grasses and Weeds In Turf and woody Ornamental shrubs, vines, and trees.

EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A

(If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> • Hold eyes open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.</p>	

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with any- thing that has been treated, such as plants, soil, or water, is coveralls over short-sleeved shirt and short pants, waterproof gloves, chemical-resistant footwear plus socks and chemical-resistant headgear for overhead exposure.

Causes substantial but temporary eye injury. Harmful if absorbed through skin. Harmful if inhaled. Do not get in eyes or clothing. Wear appropriate protective eyewear. Avoid contact with skin. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

- Shoes plus socks
- Protective eyewear
- Waterproof gloves

User Safety Recommendations	
Users should:	<ul style="list-style-type: none"> Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. Oxa-Pro 2G ® should not be used in areas where irrigation or rainfall results in the direct or indirect (via ditches and canals) contamination of surface waters through dissolved runoff or erosion of soil particles to which oxadiazon is absorbed.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

Do not apply THIS PRODUCT through any type of irrigation system.

The Andersons Oxa-Pro 2G is a selective, preemergent herbicide for control of many annual grasses and broadleaf weeds in turf and on a wide variety of woody ornamental shrubs, vines, and trees. The Andersons Oxa-Pro 2G controls weeds by killing the young weed seedlings as they come in contact with the herbicide during germination.

Apply The Andersons Oxa-Pro 2G with a properly calibrated granular applicator that will apply the granules uniformly (uniform application is essential for satisfactory weed control). Calibrate the applicator prior to use, according to the manufacturer's directions, and check frequently during the application to be certain that the equipment is working properly.

Do not allow domestic animals to graze treated areas. Avoid contamination of water intended for irrigation and domestic use.

For use in established perennial Bluegrass, Bentgrass, Bermudagrass, Buffalograss, Perennial Ryegrass, St. Augustinegrass, Seashore Paspalum*, Tall Fescue, and Zoysia turf; such as, fairways, parks, golf courses, and lawns. Rake turf area free of leaves, mowing if necessary, so granules reach soil surface. Measure area to be treated and uniformly apply required amount, using a fertilizer spreader. Cross-treat with one-half required amount in each direction will give best results. Do not apply to wet turf. If rain is not expected shortly after application, irrigating turf thoroughly will increase effectiveness.

- Do not apply to Bentgrass turf moved at less than 3/8 inch.
- Do not apply this product to newly seeded areas.
- Do not apply to Dichondra or Centipedegrass.
- Do not apply this product to putting greens or tees because of the varying cultural practices used.
- Do not apply more than 4 lbs. of active ingredient per acre per application.
- Do not apply to wet turf.
- Do not apply more than 200 lbs. of product (4 lbs. of ai) per acre per application.
- Not for use in turfgrass on Residential properties.
- The maximum application rate allowed per year is 300 lbs. of product per acre per year (equivalent to 6 lbs. of ai per acre per year), except in areas where there is heavy weed infestation. In areas of heavy weed infestation, the maximum application rate is 400 lbs. of product per acre per year (equivalent to 8 lbs. of ai per acre per year).
- For St. Augustinegrass and Bentgrass turf, do not exceed the rate of 150 lbs. per acre per year (3 lbs of ai) of this product.
- Do not overseed fairways before 60 days after an application of this product. Do not use where Poa Trivialis is present in overseed mix.

- This product may cause discoloration of Bentgrass, Bermudagrass and St. Augustinegrass. The discoloration is temporary and normally outgrown within 2 to 3 weeks.

- Avoid contact with flowers and shrubs except as discussed elsewhere on this label.
- Seeding into treated areas should be delayed until 4 months after treatment except as noted below.
- When making applications to Tropic Lalo, applications should be made 0-12 days after sprigging.

PRE-EMERGENT WEED CONTROL			
	AMOUNT OF PRODUCT		
WEEDS CONTROLLED	PER ACRE	PER 1000 SQ. FT.	APPLICATION INSTRUCTIONS
Crabgrass (<i>Digitaria spp.</i>) Goosegrass (<i>Eleusine indica</i>) *Field Sandbur (<i>Cenchrus incertus</i>) *Annual Sedge (<i>Cyperus compressus</i>)	100 - 200 lbs.	2 ¼ - 4 ½ lbs. Use the higher rate with expected heavier weed pressure	Apply in late winter or early spring prior to seed germination. Application made after March or April may not be fully effective.
*Annual Bluegrass (<i>Poa Annua spp.</i>) (See note below)	100-200 lbs.	2 ¼ - 4 ½ lbs. Use the higher rate with expected heavier weed pressure	Apply in late summer to early fall prior to seed germination. The Andersons Oxa-Pro 26 does not control <i>Poa annua</i> that has germinated prior to the application.
Virginia Buttonweed (<i>Diodia virginiana</i>) Pigweed (<i>Amaranthus spp.</i>) Florida Pusley (<i>Richardia scabra</i>) Oxalis (<i>Oxalis spp.</i>) Stinging Nettle (<i>Urtica dioica</i>) Carpetgrass (<i>Axonopus officinis</i>)	200 lbs.	4 ½ lbs.	Apply in early spring prior to seed germination.

NOTE: In many areas, annual bluegrass exists as at least two biotypes or subspecies of *Poa annua*. That true annual biotype is *Poa annua* spp. which germinates every year from seed, while the biotype that behaves like a perennial is *Poa annua* spp. reptans. This perennial bluegrass can survive through the summer and established plants will not be controlled by this product. Consult the extension service or University weed specialists in your area for more information.

*Not currently registered in California.

FALL APPLICATION FOR SPRING WEED CONTROL IN COOL SEASON TURF*			
	AMOUNT OF THIS PRODUCT		
WEEDS CONTROLLED	PER ACRE	PER 1000 SQ. FT.	APPLICATION INSTRUCTIONS
Crabgrass, Goosegrass and other annual weeds	150 - 200 lbs.	3 ¼ - 4½ lbs. Use the higher rate with expected heavier weed pressure	For use in Bluegrass, Bentgrass, Perennial Ryegrass and Tall Fescue turf. Make application in the fall after turf has become dormant. For late, fall applications, seeding into treated areas must be delayed until 4 months after green up.
ANNUAL WEED CONTROL DURING BERMUDAGRASS, ZOYSIAGRASS, SEASHORE PASPALUM, AND TROPIC LALO (Paspalum Hieronymii) ESTABLISHMENT FROM SPRIGS*			
	AMOUNT OF OXA-PRO 2G		
WEEDS CONTROLLED	PER ACRE	PER 1000 SQ. FT.	APPLICATION INSTRUCTIONS
Crabgrass, Goosegrass, Annual Sedge, and other annual weeds	100 to 150 lbs.	2 ¼ to 3 ¾ lbs. Use the higher rate with expected heavier weed pressure	For Bermudagrass , make application immediately prior to or immediately after sprigging. For Seashore Paspalum , make application 10 to 14 days after sprigging. Applications made close to the time of sprigging may cause objectionable Seashore Paspalum injury. Disturbance of the soil surface after The Andersons Oxa-Pro 2G application may disrupt the herbicide barrier and reduce effectiveness. Irrigate immediately after application for best results.
RESTRICTIONS			
• For St. Augustinegrass and Bentgrass turf, do not exceed the rate of 150 pounds per acre (3 pounds active ingredient) of this product. Do not apply to Bentgrass turf mowed at less than ¾ inch. The maximum number of applications that may be made per year are two. The maximum single application rate is 4 lbs. ai per acre per year. A subsequent application can be made 4 months after the previous treatment.			

The Andersons Oxa-Pro 2G may cause discoloration of Bentgrass, Bermudagrass and St. Augustinegrass. This discoloration is temporary and normally outgrown within 2 to 3 weeks.

Avoid contact with flowers and shrubs except as directed elsewhere on this label. Do not apply The Andersons Oxa-Pro 2G to newly seeded areas. Seeding into treated areas must be delayed until 4 months after treatment except as noted above. Do not apply to Dichondra or centipedegrass. The Andersons Oxa-Pro 2G must not be used on putting greens or tees because of the varying cultural practices used.

When making applications to TropicLalo, applications must be made 0-12 days after sprigging. Do not use in areas where endangered Hawaiian Goose or nene (*Branta sandvicensis*), Hawaiian duck (*Anas wyvilliana*), Hawaiian Stilt (*Himantopus mexicanus knudsem*), and Hawaiian Moorhen (*Gallinula chloropus sandvicensis*) may feed.

Apply The Andersons Oxa-Pro 2G anytime during the year, prior to weed seed germination, for weed control. The Andersons Oxa-Pro 2G can be applied to both newly transplanted and established ornamentals. Application can be made to actively growing or dormant ornamentals. Rainfall or overhead irrigation after application of The Andersons Oxa-Pro 2G will improve weed control activity. The Andersons Oxa-Pro 2G should not be mixed into the soil. Cultivating or disturbing the soil may reduce the weed control activity of The Andersons Oxa-Pro 2G. **Do not** apply to wet foliage or under conditions in which granules will collect on leaves. Do not apply in greenhouses as plant injury may result.

Ground Application - Apply The Andersons Oxa-Pro 2G broadcast to ornamentals at 100 to 200 pounds of product per acre (or 2½ to 4½ pounds per 1,000 sq. ft.) depending on the weeds to be controlled, as listed below. Remove existing weed growth before application.

ANNUAL BROADLEAF WEEDS		ANNUAL GRASSES	
Bittercress	(<i>Cardamine</i> spp.)	Annual Bluegrass	(<i>Poa annua</i>)
*Bristly Oxtongue	(<i>Picris echioides</i>)	*Annual Sedge	(<i>Cyperus compressus</i>)
Carpetweed	(<i>Mo/lugo vertici/lata</i>)	*Barnyard grass	(<i>Echinochloa crus-ga/II</i>)
*Cheeseweed	(<i>Malva parviflora</i>)	*Bristly Foxtail	(<i>Setaria verticillata</i>)
Common Groundsel	(<i>Senecio vulgaris</i>)	Crabgrass	(<i>Digitaria</i> spp.)
Common Purslane	(<i>Portulaca oleracea</i>)	Fall Panicum	(<i>Panicum dichotomiflorum</i>)
*Evening Primrose	(<i>Oenothera</i> spp.)	Green Foxtail	(<i>Setaria viridis</i>)
*Fiddleneck	(<i>Amsinckia intermedia</i>)	Goosegrass	(<i>Elyusine indica</i>)
*Fireweed	(<i>Epilobium angustifolium</i>)	**Jungle Ricegrass	(<i>Echinochloa colonum</i>)
*Galinisoga	(<i>Galinisoga spp.</i>)	**Panama Paspalum	(<i>Paspalum fimbriaatum</i>)
*Garden Spurge	(<i>Euphorbia hirta</i>)	*Rigput Bromegrass	(<i>Bromus catharticus</i>)
Golden Ragwort	(<i>Senecio aureus</i>)	*Wild Oats	(<i>Avena fatua</i>)
Lambsquarters	(<i>Chenopodium album</i>)		

<p>**Liverwort 'Niruri Pennsylvania Smartweed Pettly Spurge **Prostrate Spurge Redroot Pigweed 'Shepherdspurse 'Sensitive Plant Sow Thistle Speedwell 'Spiny Amaranth Spotted Gatsear 'Swinecress Yellow Woodsorrel</p>	<p>(<i>Marchantia</i> spp.) (<i>Phy/lanthus niruri</i>) (<i>Polygonum pensylvanicum</i>) (<i>Euphorbia pep/us</i>) (<i>Euphorbia supina</i>) (<i>Amaranthus retroflexus</i>) (<i>Capse/la bursa-pastoris</i>) (<i>Mimosa predica</i>) (<i>Sonchus oleraceus</i>) (<i>Veronica spp</i>) (<i>Amaranthus spinosus</i>) (<i>Hypochaeris radicata</i>) (<i>Coronopus didymus</i>) (<i>Oxalis stricta</i>)</p>	<p>*Use 150 to 200 pounds (3.0 to 4.0 pounds of active ingredient) per acre for less susceptible weeds. **Use 200 pounds (4.0 pounds of active ingredient) per acre for less susceptible weeds. ' Use not permitted in California</p>
---	--	---

Suggested Spreader Settings
 These suggested spreader settings are not intended to replace calibration.
 Please calibrate your spreader before applying product.

50 lbs. treats 22,200 sq. ft. (0.51 acre) at the 98 lbs. product/acre - LIGHT RATE (2.25 lbs/1,000 sq. ft.)	50 lbs. treats 14,700 sq. ft. (0.34 acre) at the 147 lbs. product/acre - LOW RATE (3.38 lbs/1,000 sq. ft.)
50 lbs. treats 13,333 sq. ft. (0.31 acre) at the 163 lbs. product/acre - MEDIUM RATE (3.75 lbs/1,000 sq. ft.)	50 lbs. treats 11,100 sq. ft. (0.26 acre) at the 196 lbs. product/acre - HIGH RATE (4.5 lbs/1,000 sq. ft.)

SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS			
			LIGHT RATE 1.96 lb. ai/A	LOW RATE 2.95 lb. ai/A	MEDIUM RATE 3.27 lb. ai/A	HIGH RATE 3.92 lb. ai/A
Andersons Model 2000 / AccuPro 2000 (cone 9); Adersons 2000 SR / SR-2000 (cone 9); Andersons LCO-1000	3 mph	10 feet	1 1/4	1 3/4	1 3/4	1 3/4
Scotts R-8, R-8A (cone 9)	3 mph	10 feet	1 3/4	1 1/4	1 3/4	1 1/4
LESCO 020093 (0.25)	3 mph	7 feet	1 1/2	1 1/4	1 1/2	1 3/4
LESCO 705698 (0.25)	3 mph	7 feet	9	10 1/2	11	11 1/2
The Andersons SSD/SS-2 Drop	3 mph	overlap wheels	3	3 1/4	3 1/2	4

NOTE: These settings are for initial calibration only. It is strongly advised that your equipment be calibrated prior to actual application, using these settings as a starting point.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If not available, then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE: Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. To the extent consistent with applicable law, by using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION OF LIABILITY: Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS: THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION. IMPORTANT: These settings are only approximate.

ATTENTION: This product contains a chemical known to the State of California to cause cancer.

Manufactured by:
The Andersons, Inc.
P.O. Box 119
Maumee, OH 43537
www.andersonsplantnutrient.com



Oxa-Pro is a trademark of The Andersons, Inc.

Net Weight 50 lbs. (22.68 kg)

10007945
K22